



Noggin rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A18480	Rabbit	1 mg/ml	kD
Applications	IHC,ELISA		
Reactivity	Human,Mouse,Rat		
Dilution	IHC 1:50-200, ELISA 1:10000-20000		
Storage	-20°C/1 year		
Specificity	The antibody detects endogenous Noggin		
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human NOG. AA range:21-70		
Uniprot No	Q13253		
Alternative names	Noggin		
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype			
Conjugation			
Background	noggin(NOG) Homo sapiens The secreted polypeptide, encoded by this gene, binds and inactivates members of the transforming growth factor-beta (TGF-beta) superfamily signaling proteins, such as bone morphogenetic protein-4 (BMP4). By diffusing throu		
Other	Gene_name: NOG ; Protein_name: Noggin; Expression: Placenta,Prostate,Temporal cortex,		

Product Images

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:



H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.