



## PJVK rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A19878	Rabbit	1 mg/ml	38 kD
Applications	IF,ELISA		
Reactivity	Human		
Dilution	IF: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.		
Storage	-20°C/1 year		
Specificity	PJCG2 Polyclonal Antibody detects endogenous levels of PJCG2 protein.		
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
Immunogen	The antiserum was produced against synthesized peptide derived from human PJCG2. AA range:140-189		
Uniprot No	Q8WZ85		
Alternative names			
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype			
Conjugation			
Background	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from sing		
Other	Gene_name: PJCG2 ; Protein_name: Olfactory-like receptor PJCG2; Expression:		

### Product Images

#### Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation  
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

#### Species Cross-Reactivity Key:



H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster  
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All  
Species Expected

**Trademarks**

*All product names and trademarks are the property of their respective owners.*

**Regulatory Disclaimer**

*For life science research only. Not for use in diagnostic procedures.*

---

**Contact and Support:**

*To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).*

*To call, write, fax, or email us, please visit [www.aabsci.com](http://www.aabsci.com), contact information will be displayed.*