



PNDC1 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A20042	Rabbit	1 mg/ml	57 kD
Applications	WB		
Reactivity	Human, Mouse,Rat		
Dilution	WB 1:500-2000		
Storage	-20°C/1 year		
Specificity	This antibody detects endogenous levels of PNDC1 at Human/Mouse/Rat		
Source / Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.		
Immunogen	Synthesized peptide derived from human PNDC1		
Uniprot No	Q6XQN6		
Alternative names			
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype			
Conjugation			
Background	Nicotinic acid (NA; niacin) is converted by nicotinic acid phosphoribosyltransferase (NAPRT; EC 2.4.2.11) to NA mononucleotide (NaMN), which is then converted to NA adenine dinucleotide (NaAD), and finally to nicotinamide adenine dinucleotide (NAD), which		
Other	Gene_name: NAPRT1 FHIP ; Protein_name: PNDC1; Expression:		
Product Images			

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All



Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.