



PROL4 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A20222	Rabbit	1 mg/ml	14 kD
Applications	IHC, ELISA		
Reactivity	Human, Mouse, Rat		
Dilution	IHC 1:50-200, ELISA(peptide)1:5000-20000		
Storage	-20°C/1 year		
Specificity	This antibody detects endogenous levels of human Prokineticin-1		
Source / Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.		
Immunogen	Synthesized peptide derived from human Prokineticin-1		
Uniprot No	P58294		
Alternative names	Prokineticin-1 (Endocrine-gland-derived vascular endothelial growth factor) (EG-VEGF) (Mambakine)		
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype			
Conjugation			
Background	prokineticin 1(PROK1) Homo sapiens The protein encoded by this gene induces proliferation, migration, and fenestration (the formation of membrane discontinuities) in capillary endothelial cells derived from endocrine glands. It has little or no effect		
Other	Gene_name: PROK1 UNQ600/PRO1186 ; Protein_name: Prokineticin-1 ; Expression: Testis,		

Product Images

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:



H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.