



β II Tubulin rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A23883	Mouse	1 mg/ml	50 kD

Applications	WB,IHC,IP
Reactivity	Mouse,Rat,(H)
Dilution	WB: 1:50000-100000 IP:1:200 IHC 1:50-300
Storage	-20°C/1 year
Specificity	The antibody detects endogenous β II tubulin protein.
Source / Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Immunogen	Synthetic Peptide of β II tubulin
Uniprot No	Q13885
Alternative names	Tubulin beta-2A chain
Form	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Clonality	Polyclonal
Isotype	
Conjugation	
Background	tubulin beta 2A class IIa(TUBB2A) Homo sapiens Microtubules, key participants in processes such as mitosis and intracellular transport, are composed of heterodimers of alpha- and beta-tubulins. The protein encoded by this gene is a beta-tubulin.
Other	Gene_name: TUBB2A ; Protein_name: Tubulin beta-2A chain; Expression: Brain,Cajal-Retzius cell,Eye,Fetal brain cortex,Skin,

Product Images

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:



H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.