



GST mouse mAb(Q16) antibody

| Catalog No : | Source: | Concentration : | Mol.Wt. (kD): |
|-----------------------|---|-----------------|---------------|
| A42012 | Mouse | | |
| Applications | WB | | |
| Reactivity | Species independent | | |
| Dilution | WB: 1:2000 -5000 | | |
| Storage | -20°C/1 year | | |
| Specificity | The antibody detects C-terminal, internal, and N-terminal GST fusion proteins. | | |
| Source / Purification | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. | | |
| Immunogen | Recombinant Protein of GST | | |
| Uniprot No | | | |
| Alternative names | | | |
| Form | PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol. | | |
| Clonality | Polyclonal | | |
| Isotype | | | |
| Conjugation | | | |
| Background | GST (Glutathione S-Transferase) is a widely used protein tag encoded by the Schistosoma japonicum. GST provides both an easily detectable tag and a simple purification process with little effect on the biological function of the protein of interest. Antib | | |
| Other | | | |
| Product Images | | | |

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:



H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.