



Cystatin C mouse mAb(7F11) antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A13345	Mouse	1 mg/ml	14 kD
Applications	IF, WB, IHC, ELISA		
Reactivity	Human		
Dilution	IF: 1:50-200 WB 1:1000-2000, IHC 1:100-200		
Storage	-20°C/1 year		
Specificity	Cystatin C protein detects endogenous levels of CST3		
Source / Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.		
Immunogen	Recombinant Protein of Cystatin C of CST3		
Uniprot No	P01034		
Alternative names	CST3		
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
Clonality	Monoclonal		
Isotype			
Conjugation			
Background	cystatin C(CST3) Homo sapiens The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibit		
Other	Gene_name: CST3 ; Protein_name: CST3; Expression: Brain, Leukocyte, Placenta, Synovial cell,		

Product Images

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster



X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.