



IL-17C rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A16191	Rabbit	1 mg/ml	kD
Applications	IHC,ELISA		
Reactivity	Human		
Dilution	IHC 1:50-200, ELISA 1:10000-20000		
Storage	-20°C/1 year		
Specificity	The antibody detects endogenous IL-17C		
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
Immunogen	Synthetic peptide from human protein at AA range: 81-130		
Uniprot No	Q9P0M4		
Alternative names	Interleukin-17C (IL-17C) (Cytokine CX2)		
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype			
Conjugation			
Background	interleukin 17C(IL17C) Homo sapiens The protein encoded by this gene is a T cell-derived cytokine that shares the sequence similarity with IL17. This cytokine was reported to stimulate the release of tumor necrosis factor alpha and interleukin 1 beta		
Other	Gene_name: IL17C UNQ561/PRO1122 ; Protein_name: Interleukin-17C (IL-17C) (Cytokine CX2); Expression: Lung,		

Product Images

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster



X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.