



IL-17D rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A16192	Rabbit	1 mg/ml	kD

Applications	IHC,ELISA
Reactivity	Human
Dilution	IHC 1:50-200, ELISA 1:10000-20000
Storage	-20°C/1 year
Specificity	The antibody detects endogenous IL-17D
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Immunogen	Synthetic peptide from human protein at AA range: 141-190
Uniprot No	Q8TAD2
Alternative names	Interleukin-17D (IL-17D) (Interleukin-27) (IL-27)
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Clonality	Polyclonal
Isotype	
Conjugation	
Background	interleukin 17D(IL17D) Homo sapiens The protein encoded by this gene is a cytokine that shares the sequence similarity with IL17. The treatment of endothelial cells with this cytokine has been shown to stimulate the production of other cytokines inclu
Other	Gene_name: IL17D IL27 UNQ3096/PRO21175 ; Protein_name: Interleukin-17D (IL-17D) (Interleukin-27) (IL-27); Expression: Brain,

Product Images

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster



X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.